



Age-related differences in affective responses to and memory for emotions conveyed by music: a cross-sectional study

Sandrine Vieillard^{1*} and Anne-Laure Gilet²

¹ Laboratoire de Psychologie (EA 3188), Psychology, Université de Franche-Comté, Besançon, France

² Laboratoire de Psychologie des Pays de la Loire (EA 4638), Université de Nantes, Nantes, France

Edited by:

Petri Laukka, Stockholm University, Sweden

Reviewed by:

Jonna K. Vuoskoski, University of Oxford, UK

Michaela Riediger, Max Planck Institute for Human Development, Germany

*Correspondence:

Sandrine Vieillard, Laboratoire de Psychologie (EA 3188), Université de Franche-Comté, 30 Rue Mégevand, 25030 Besançon, France
e-mail: sandrine.vieillard@univ-fcomte.fr

There is mounting evidence that aging is associated with the maintenance of positive affect and the decrease of negative affect to ensure emotion regulation goals. Previous empirical studies have primarily focused on a visual or autobiographical form of emotion communication. To date, little investigation has been done on musical emotions. The few studies that have addressed aging and emotions in music were mainly interested in emotion recognition, thus leaving unexplored the question of how aging may influence emotional responses to and memory for emotions conveyed by music. In the present study, eighteen older (60–84 years) and eighteen younger (19–24 years) listeners were asked to evaluate the strength of their experienced emotion on happy, peaceful, sad, and scary musical excerpts (Vieillard et al., 2008) while facial muscle activity was recorded. Participants then performed an incidental recognition task followed by a task in which they judged to what extent they experienced happiness, peacefulness, sadness, and fear when listening to music. Compared to younger adults, older adults (a) reported a stronger emotional reactivity for happiness than other emotion categories, (b) showed an increased zygomatic activity for scary stimuli, (c) were more likely to falsely recognize happy music, and (d) showed a decrease in their responsiveness to sad and scary music. These results are in line with previous findings and extend them to emotion experience and memory recognition, corroborating the view of age-related changes in emotional responses to music in a positive direction away from negativity.

Keywords: aging, musical emotions, emotional responses, facial muscle activity, incidental recognition, positivity effect

INTRODUCTION

Research on age differences in emotion processing has been mostly restricted to visual stimuli (e.g., facial expression, video, words, and pictures) but a growing body of research converges in indicating that music also serves as a powerful emotional trigger. For example, Blood et al. (1999) have shown that classical musical stimuli which are selected to elicit intensely pleasant emotional responses engage neural networks that are implicated in reward. Neuropsychological studies have also demonstrated that the amygdala is recruited when processing scary music (e.g., Gosselin et al., 2005, 2007). Physiological data have also put in evidence that music is a strong emotion inducer (e.g., Khalfa et al., 2002). Furthermore, music has the clear advantage of maintaining attention toward the emotions conveyed because it does not allow perceptual attention to be redirected (except if the listener takes off his/her headphones). For these reasons, music appears as a viable method to test age-related changes in emotion processing.

Among the past studies that have addressed the question of age-related changes in musical emotion processing, most of them have focused on the people's ability to recognize musical emotions (Allen and Brosigole, 1993; Laukka and Juslin, 2007; Drapeau et al., 2009; van Tricht et al., 2010; Lima and Castro, 2011; Vieillard et al., 2012). For instance, Drapeau et al.

(2009) compared healthy elderly adults and elderly adults with Alzheimer's disease in their ability to rate the extent to which the selected musical stimuli communicating happiness, peacefulness, sadness and fear (Vieillard et al., 2008) expressed each of these four emotions. Their findings showed that recognition performances of healthy older adults were relatively preserved with the highest recognition accuracy for happy stimuli. Laukka and Juslin (2007) compared young and older adults' ability to recognize anger, fear, happiness, sadness, and neutrality in short melodies performed on an electric guitar with different degrees of expressivity. In their study, the participants judged the emotional expression of each stimulus in a forced choice task comprised of anger, fear, happiness, sadness, neutral, and "other emotion" alternatives. Compared to young adults, older adults were less accurate in recognizing negative emotions such as sadness and fear, but their ability to recognize other emotion categories was spared. More recently, Lima and Castro (2011) went one step further by examining age-related changes in emotion recognition among three age groups of participants (i.e., young, middle-aged, and old). Selecting the same musical excerpts as those used in Drapeau et al. (2009)'s study, the authors asked the participants to judge the emotional intensity perceived for each stimulus on four rating scales (i.e., happiness, sadness, peacefulness, and fear)

presented simultaneously. Consistent with previous findings, the authors found an emotion-specific age-related change characterized by a stable recognition of happiness and peacefulness categories but a gradual decline in responsiveness to sad and scary music from young adulthood to older age.

As an alternative to the hypotheses of age-related differences in cognitive functioning¹ or hearing loss², the above findings have been interpreted as being the result of a combination of age-related changes in brain structure and functioning and in motivational goals. On the one hand, empirical evidence suggest that the observed decline found in older adults in negative emotion recognition may be the result of a linear reduction in the volume of the amygdala (e.g., Zimmerman et al., 2006) and/or of a decrease of the reactivity to negative information in the amygdala (e.g., Mather et al., 2004). On the other hand, Socioemotional Selectivity Theory (Carstensen et al., 1999; Carstensen, 2006) suggests that the decline in the recognition of negative emotions would reflect a motivational shift toward emotionally meaningful goals due to an increased awareness of the limited perspective of time. This so-called “positivity effect” which refers to all combinations of *enhanced* processing of positive information and *reduced* processing of negative information, has been thought of as an emotion regulation strategy to preserve high levels of well-being in later life. In a recent literature review, Reed and Carstensen (2012) showed that this positivity effect (1) requires cognitive resources (e.g., Mather and Knight, 2005), (2) is sensitive to the experimental context (Kensinger et al., 2002; Grühn et al., 2005), and (3) is adaptive, i.e., it emerges when emotional well-being is prioritized. Consequently, the authors claimed that the positivity effect would represent a controlled shift in attentional resources rather than an *automatic* process associated with the neuronal degeneration in brain regions. Such view is compatible with the idea that the positivity effect may have a cognitive counterpart.

Recently, Vieillard et al. (2012) conducted a study in order to test for age-related differences in the psychological structure of musical emotions, and to assess whether these changes may be associated with a decrease in emotional complexity. In this research, younger and older participants were presented with musical excerpts conveying different emotions such as happiness, peacefulness, sadness, and fear (Vieillard et al., 2008). Participants were asked to perform an emotional judgment task using different rating scales (i.e., valence, hedonic value, arousal, and liking) as well as a free categorization task in which they freely created emotional categories based on the perceived acoustical cues. Findings showed age-related differences characterized by a reduced processing of arousal for scary music, an increased focus on happy music, and an emotional dedifferentiation corresponding to a decrease in differentiation between the arousal and valence dimensions. Such results have been explained within the framework of the Dynamic Integration Theory (Labouvie-Vief, 2009; Labouvie-Vief et al., 2010) postulating that the degradation of emotional complexity would be the cognitive counterpart of the

older adults’ attempt to maximize positive affects and minimize negative affect in order to preserve well-being.

In short, the studies reviewed above show converging evidence for a positivity effect in how emotions in music are perceived, categorized, and recognized with advancing age. However, several questions remain open. First, little is known about age-related changes regarding the emotions *experienced* while listening to music. Previous findings have suggested that participants were more accurate in their judgment of intended emotions in musical excerpts when focusing on their own emotional experience (Vieillard et al., 2008). One of the goals of the present study was thus to investigate the influence of aging on emotion processing while being personally engaged in musical listening. This is an important question because it has been suggested that emphasizing on emotion rather than on knowledge may be more meaningful to older adults (e.g., Mikels et al., 2010). Second, as far as we know, the possibility of age-related changes in memory recognition for positive musical stimuli has yet to be examined. Past research showed that age-related changes in emotional goals influence memory. A positivity effect in memory recognition tasks has already been shown in older adults for affective pictures (e.g., Charles et al., 2003; Mather and Carstensen, 2005) and for words (e.g., Kensinger, 2008). The main explanation was that memory can work as an elaborative process to regulate emotions such that the older adults’ goal to maintain well-being would influence mental constructions of the past and thus lead to a positivity effect in the way they remember events. In line with this hypothesis, it has been showed that memory for negative pictures decreased in older adults both in recall and recognition tasks (e.g., Charles et al., 2003). Kensinger (2008) found a positivity effect in older adults for non-arousing words, explaining this as an age-related difference in the way positive information was primarily processed as a function of differences in motivational goals at each age level. To date, the question remains open whether age-related differences in the elaborative processing of memory may be observed in a non-verbal channel of emotion communication such as music. Third, to our knowledge, no previous study has investigated age differences in facial muscle activity when listening to music. Past studies examining age-related differences in facial expressiveness have found that young and older adults express similar patterns of facial responding to visual stimuli such as emotional scenes, objects or faces (Reminger et al., 2000; Smith et al., 2005; Bailey et al., 2009). However, older adults compared to younger adults may exhibit diminished reactivity in facial expressiveness (Smith et al., 2005; Burriss et al., 2007). The reduction of facial expressiveness in the elderly has been thought to be a possible consequence of general physiological losses in the nervous system. However, another explanation suggests that the reduction of facial expressiveness may reflect an attempt to regulate emotion since facial expression may be motivationally driven (Smith et al., 2005). Given such emotion regulation hypothesis and in line with the embodiment theory of emotion (Niedenthal et al., 2005), one can imagine that facial expressions may not only help to down-regulate emotion (by displaying less facial expressions), but also allow to modify the emotional reaction (by displaying a facial expression contrary to what one feels). In this perspective, facial electromyogram (EMG) appears as an interesting indicator

¹ Emotion categories that were less accurately recognized by older adults were not those known to be the hardest to recognize.

² Age groups did not differ or only differed marginally on self-reported hearing loss.

of whether there is congruence between facial expressivity and musical emotion in both young and older adults, or do older adults express positive facial expression as a means to counteract negative emotions.

There is an agreement to consider that emotional responding is a multi-component process, giving rise to affective experiences, physiological adjustments and expressive behaviors (e.g., Scherer, 2005). These various aspects of emotion may be differentially influenced by age. Therefore, a unifying view of these changes is necessary to give more insight into the lifespan developmental course of emotion, particularly in the musical domain in which this topic has remained unexplored. To this end, we focused on different indexes of emotional response to music, namely subjective experience, emotion expression and memory recognition for musical excerpts that conveyed different emotions.

Finally, although the positivity effect has been observed across a number of experimental paradigms such as dot-probe tasks (Charles et al., 2003), eye-tracking paradigms (Isaacowitz et al., 2006a,b), working memory (Mikels et al., 2005), memory recognition and free recall tasks (Charles et al., 2003), and across a variety of stimuli (e.g., pictures, word lists, facial expressions), the robustness of this phenomenon has been mainly demonstrated through the visual channel of emotion communication. Consequently, research regarding the effect of age on emotion processing in music is needed to test for the generalizability of the positivity effect.

CURRENT RESEARCH

The present study was designed to further extend previous studies and expand experimental designs to the domain of music. Our aim was twofold: first, to investigate the effects of aging on the emotion felt when listening to music and second, to address age-related differences on memory recognition for musical excerpts as a function of their intended emotion. To this end, we used a set of musical stimuli expressing happiness, peacefulness, sadness, and fear which were all controlled for valence and arousal (Vieillard et al., 2008). We used a rating task focused on the emotion experienced by the participants rather than on the emotion recognized by the participants. In order to address more extensively the effect of aging on the experienced emotion, we designed the experiment so that the subjective report of the intensity of the emotion felt was coupled with a recording of participants' facial expressions. These particular indexes were chosen since past research has shown that facial expressions, measured by the corrugator (i.e., frowning) and zygomatic (i.e., smiling) muscle activity, were mostly related to the valence in music: positive emotions generally lead to increased zygomatic activity, while negative ones were associated more with increased corrugator activity (e.g., Witvliet and Vrana, 2007; Khalifa et al., 2008). Because it has been suggested that facial EMG may also be voluntarily modulated to serve emotion regulation goal (Smith et al., 2005), we used this index to examine to what extent and how older adults show positive or negative facial expression as a function of musical emotions.

Based on the hypothesis postulating a motivated attention toward positivity with advancing age, and given the view that situations relevant to a person's motivational goals may elicit more intense emotional experience (e.g., Charles and Piazza,

2007), we expected that older adults, compared to younger adults, would judge their feeling as more intense when listening to positive musical excerpts (especially happy music that is more arousing than peaceful one) than when listening to negative musical excerpts. Since no age-related changes in facial expressivity were found in previous research (e.g., Levenson et al., 1991; Tsai et al., 2000; Magai et al., 2006), we also expected older adults to be spared in their facial expressions (i.e., corrugators and zygomatic muscle activity). More specifically, if older adults have spontaneous facial activity, it is expected that they would display a greater zygomatic activity for positive music in comparison with their younger counterparts. At the same time, it is also expected that older adults would display voluntary facial expression as a tool to manage emotion. In this hypothesis, older adults would show reduced expressivity or incongruent expression, in particular in response to negativity. Moreover, in view of the scarce data available on the influence of aging on memory recognition for musical excerpts that convey different emotions, and because the memory elaborative processes for music are based on more abstract information than those involved in the memory for visual and autobiographical material, it is difficult to predict the nature of the effects likely to be observed. Consequently, we conducted an exploratory approach to test whether the positivity effect may be generalized to memory recognition for musical excerpts conveying different emotions. We expected that compared to younger adults, older adults should better recognize positive musical excerpts than negative ones.

METHOD

PARTICIPANTS

A total of 40 native French speaking volunteers (22% amateur musicians³) participated in the present study. Exclusion criteria included the presence of uncorrected hearing, medical or psychiatric antecedents, psychotic symptoms, and history of substance abuse. As a result, the data of 18 young adults (19–24 years, $M = 21$ years; 61 % females) and 18 older adults (60–84 years, $M = 66$ years; 83% females) was analyzed. Younger and older adults were recruited, respectively at the psychology department of the University of Franche-Comté and through senior social programs in Besançon. Participants did not receive financial compensation for their participation.

APPARATUS

Participants were tested individually in a quiet room at stable ambient temperature at the University. Facial muscle activity was monitored continuously during the listening and rating phases using an MP150 Biopac system (Biopac Systems, Inc., Goleta, CA) at a sampling rate of 500 Hz and processed using AcqKnowledge software. Eprime software (Schneider et al., 2002) was used for excerpts presentation and ratings recording. Musical excerpts were presented binaurally through Professional 240 Sennheiser headphones.

³The musical training was measured as the proportion of participants who received at least 3 years of formal training and who were still practicing a musical instrument without reaching professional levels.

MATERIALS

Forty short musical excerpts, computer-generated in a piano timbre and taken from Vieillard et al. (2008) set of unfamiliar musical stimuli were selected for their power to convey four distinct emotions (i.e., happiness, peacefulness, sadness, and fear). Musical excerpts were controlled for their valence (unpleasant vs. pleasant) and arousal (low vs. high). Each emotion category included ten musical excerpts that lasted an average of 10 s. The happy excerpts were written in a major mode at an average tempo of 137 Metronome Markings (MM range: 92–196), with the melodic line lying in the medium high pitch range (the pedal was not used). The peaceful excerpts were composed in a major mode, had an intermediate tempo (mean: 74 MM, range: 54–100), and were played with pedal and arpeggio accompaniment. The sad excerpts were written in a minor mode at an average slow tempo of 46 MM (range: 40–60), with the pedal. The scary excerpts were composed with minor chords on the third and sixth degree, hence implying the use of many out-of-key notes. Although most scary excerpts were regular and consonant, a few had irregular rhythms and were dissonant. Their tempo varied from 44 to 172 MM. Examples can be heard at www.brams.umontreal.ca/peretz. A previous study that was conducted to examine the effect of age on emotion perception in music demonstrated that older listeners successfully distinguished happiness, peacefulness, sadness, and fear conveyed by these musical excerpts (Vieillard et al., 2012). In a study phase described below, participants were presented with 20 musical excerpts (i.e., 5 happy, 5 peaceful, 5 sad, and 5 scary) and were instructed to indicate what they experienced in terms of Emotional Intensity. The 20 remaining musical excerpts were then used as lures in the incidental recognition task.

PROCEDURE

The experiment was divided into two sessions separated by an interval of ~1 week. During the first session, participants completed a consent form and were asked about their age, musical listening, education level, self-reported health, visual and auditory acuity, and medical history. Auditory perception was controlled using free AudioTest software (www.cotral.com). More specifically, it was assessed by presenting pure tones at intervals between 500 and 8000 Hz to both ears through Professional 240 Sennheiser headphones. For each participant, the lowest sound pressure level at which each frequency was detected was recorded. In addition, several tests assessing general cognitive function (MMSE, Petit et al., 1998), fluid intelligence (Raven's progressive matrices, set I, Raven et al., 1998, updated 2003), and working memory (letter-number sequencing from WAIS-III, Wechsler, 2000) were administered. The first session lasted about an hour.

During the second session, physiological sensors were attached while the participants sat comfortably in a quiet room in the presence of the experimenter. To prevent participants from focusing on their facial muscles, they were informed that the electrodes placed on their face were used to record their electrodermal activity during the experiment. At the beginning of the session, two musical excerpts different from those used in the experiment were used in order to adjust the volume of the headphones

for each participant. In the study phase, two practice excerpts (1 happy and 1 peaceful) following by 20 excerpts (5 of each emotion) were then presented binaurally. After each trial, participants were asked to rate the intensity of the emotion felt using a 10-point scale ranging from 0 "weak" to 9 "strong." Facial muscle activity was also recorded. The excerpts were presented in two pseudo-randomized orders that were created to ensure that no more than two excerpts of the same emotion category were presented consecutively. Each musical excerpt was preceded and followed by two baseline periods of at least 10 s of silence.

Before the incidental recognition task, participants completed two questionnaires to assess depression (Beck et al., 1998) and anxiety (Spielberger, 1993; Beck et al., 1998). In the recognition task participants were asked to indicate whether an excerpt had been heard before ("old") or not ("new") by pressing the appropriate key. In this phase, 40 musical excerpts (i.e., 20 old excerpts and 20 new excerpts) were randomly presented.

Finally, participants were instructed to listen to the same set of 40 musical excerpts presented binaurally in a randomized order. After each musical excerpt, participants were asked to judge to what extent they experienced "happiness," "peacefulness," "sadness," or "fear" using a 10-point scale ranging from 0 "not at all" to 9 "a lot." Accordingly, each excerpt was presented four times; each presentation was associated with one of the four emotion scales. The presentation order of each musical excerpt and each rating scale was fully randomized across participants. The second session lasted about 2 h. At the end of the session, participants were fully debriefed.

DATA ACQUISITION AND TRANSFORMATION

Facial EMG activity (μ Volts) was recorded over the left corrugator and zygomatic sites, using two pairs of 8 mm Ag/AgCl shielded electrodes filled with isotonic gel. The EMG data were band-pass filtered from 100 to 500 Hz and processed with a root mean square algorithm over 20 samples (with a 100-ms window). Recording artifacts were visually identified and discarded from the sample. These corresponded to less than 0.5% of all measurements.

RESULTS

SAMPLE CHARACTERISTICS

Younger adults reported more years of education than the older adults, $t_{(34)} = -2.33, p < 0.05$. A chi-square goodness-of-fit test (χ^2) indicated no significant differences between age groups in the proportion of formal musical training of at least 3 years $\chi^2(1, N = 36) = 4.5, p > 0.05$. Age groups did not differ regarding their depression, state anxiety, or trait anxiety scores⁴.

⁴BDI-II scores indicated that 75% ($n = 27$) of the participants scored below the cut-off (score of 11) for a minimum depression symptomatology and that 19.44% ($n = 7$) of the participants scored below the cut-off (score of 19) for at least mild depression. The remaining two participants (one young and one older adult) scored between 22 and 24, which indicates moderate depression symptomatology. STAI state anxiety scores indicated that 83.33% ($n = 30$) of the participants scored below the cut-off (score of 35) for very low state anxiety, 13.89% ($n = 5$) of them scored below the cut-off (score of 45) for low state anxiety, and that 2.78% ($n = 1$) of them scored below the cut-off (score of 55) for mild state anxiety. STAI trait anxiety scores indicated that 30.56% ($n = 11$) of the participants scored below the cut-off (score of 35) for very low

457 Non-parametric Mann-Whitney *U*-test⁵ performed on the audi- 514
 458 tory thresholds (dB) did not show statistically significant differ- 515
 459 ences between younger and older adults ($U = 10.50, z = 0.77,$ 516
 460 $p = 0.44$ for 500 Hz; $U = 0, z = 0, p = 1$ for 1000 Hz; $U =$ 517
 461 $10, z = 0.84, p = 0.40$ for 2000 Hz; $U = 8, z = 1.12, p = 0.26$ 518
 462 for 4000 Hz; $U = 33, z = 0.23, p = 0.81,$ for 8000 Hz). As was 519
 463 expected, younger adults scored better than older adults on fluid 520
 464 intelligence (Raven's progressive matrices, set I, Raven et al., 1998, 521
 465 updated 2003), $t_{(34)} = -2.84, p < 0.05$. Younger adults tended 522
 466 to perform better than older adults on a working memory test 523
 467 (Digit Span from WAIS-III, Wechsler, 2000), $t_{(34)} = -1.94, p =$ 524
 468 0.06 . There were no significant age differences on self-reported 525
 469 health, $t_{(34)} = -0.81, p = 0.42$. Finally, the Mini Mental State 526
 470 Examination (MMSE; Petit et al., 1998) scores for the older adults 527
 471 suggested no apparent signs of dementia ($M = 29.7, 28-30$). 528
 472 Sample characteristics are detailed in **Table 1**. 529

473 **EMOTION INTENSITY FELT**

474 A mixed model analysis of variance was conducted on the mean 530
 475 score of the Emotional Intensity Felt with Age Group (younger 531
 476 adults, older adults) as the between-subjects factor and Intended 532
 477 Emotion (happiness, peacefulness, sadness, fear) as the within- 533
 478 subjects factor. 534

479 **Table 1 | Sample characteristics.**

	Age group	
	Young (18)	Older (18)
DEMOGRAPHIC CHARACTERISTICS		
Age (year)	19–24	61–74
Sex	61% Women	83% Women
Education (year)*	15 (3.17)	13 (1.57)
Musical training (proportion)	0.22	0
Self reported health (Max. 5)	4.44 (0.86)	4.22 (0.79)
COGNITIVE SCORES		
Advanced progressive matrices (Set 1, Max. 12)*	10.00 (1.91)	7.78 (2.71)
Letter-number sequencing (Max. 21)	11.89 (3.05)	10.17 (2.23)
MMSE (Max. 30)	–	29.7
AFFECTIVE SCORES		
Depression (BDI-II, Max 63)	8.33 (6.79)	8.67 (5.30)
State Anxiety (STAI-Y, Max 80)	30.55 (6.35)	27.83 (6.72)
Trait Anxiety (STAI-Y, Max 80)	40.83 (10.26)	40.17 (9.57)

480 *Standard deviations are listed in parentheses. *Significant difference at $p < 0.05$.*

481 state anxiety, 41.67% ($n = 15$) scored below the cut-off (score of 45) for low 535
 482 trait anxiety, and 19.44% ($n = 7$) scored below the cut-off (score of 55) for 536
 483 moderate trait anxiety. The remaining two participants (one young and one 537
 484 older adult) scored between 57 and 58, which correspond for a high anxiety 538
 485 state. 539

486 ⁵Because the audio test software did not provide absolute thresholds, we used 540
 487 the lowest sound pressure level (dB) at which each frequency was detected 541
 488 as an ordinal data. For this reason, the non-parametric Mann-Whitney test 542
 489 (independent samples) was used. Although hearing thresholds varied as a 543
 490 function of age, the Mann-Whitney test did not reveal significant differences 544
 491 between age groups probably because of the variability observed within the 545
 492 groups, in particular in the older adults. 546

Emotions (happiness, peacefulness, sadness, fear) as the within- 514
 subjects factor. 515

As illustrated in **Figure 1**, we found a significant Age Group by 516
 Intended Emotion interaction, $F_{(3, 102)} = 3.75, p < 0.05, \eta_G^2 =$ 517
 0.06^6 . In order to test our hypothesis, we computed a planned 518
 comparison between the emotion intensity felt by young adults 519
 and that felt by older adults when listening to happy music. As 520
 expected, older adults compared to young adults reported experi- 521
 encing higher emotional intensity when listening to happy music, 522
 $F_{(1, 34)} = 5.57, p < 0.05, \eta_G^2 = 0.14$. This older adults' reactivity 523
 for positivity was also confirmed by another set of planned com- 524
 parisons indicating that older adults reported experiencing higher 525
 emotional activation when listening to happy music than when 526
 listening to sad music, $F_{(1, 34)} = 7.34, p < 0.05, \eta_G^2 = 0.18,$ or 527
 scary music, $F_{(1, 34)} = 8.76, p < 0.05, \eta_G^2 = 0.20,$ while younger 528
 adults did not. No other significant effect was found. A separ- 529
 ate analysis with years of education, fluid intelligence scores, and 530
 working memory performances (i.e., factors that were found to 531
 be different between the two age groups) as covariates indicated 532
 that the Age Group by Intended Emotion interaction remained 533
 significant, $F_{(3, 93)} = 3.14, p < 0.05, \eta_G^2 = 0.05$. 534

535 **FACIAL MUSCLE ACTIVITY**

536 Facial EMG responses were calculated as the difference between 537
 the signal (Area under the curve, $\mu V \cdot sec$) over the time course 538
 of the musical excerpt and a baseline EMG level measured from 539
 1s prior to the onset of the excerpt (time -1 to 0 s) to the begin- 540
 ning of the excerpt. The area under the curve was extracted 541
 within these two time windows and was averaged for each con- 542
 dition and for each participant. Two participants were excluded 543
 (1 younger and 1 older adults) from the initial sample due to 544
 technical problems. Analyses were then conducted on seventeen 545
 younger and seventeen older adults. The Shapiro-Wilk normality 546
 test reached significance for the sample set of EMG data mean- 547
 ing that the assumption of normality has to be rejected. As a 548
 result, statistical analyses were performed using non-parametric 549
 tests. First, the differences between age groups were tested sep- 550
 arately for zygomatic and corrugator using the Mann-Whitney 551
U-test. Results showed a significant effect of Age Group both 552
 for zygomatic ($U = 42, z = 3.51, p < 0.001$) and for corrugator 553
 muscle ($U = 83, z = 2.10, p < 0.05$) indicating that facial activ- 554
 ity was more important in older adults than in their younger 555
 counterparts. Friedman repeated measures analyses of variance 556
 (RM-ANOVA) were conducted separately on zygomatic and on 557
 corrugator activity to test the effect of Intended Emotion factor 558
 for each Age Group. For zygomatic muscle, data revealed a signifi- 559
 cant effect of Intended Emotion in older adults, $\chi^2 = 12.46, df =$ 560
 $3, p < 0.05,$ but not in young adults, $\chi^2 = 2.01, df = 3, p = 0.57$. 561
 As shown in **Figure 1**, older adults showed an increased zyo- 562
 gomatic activity, in particular for scary music. Regarding corrugator 563
 activity, older adults showed an increased activity, in particular for 564
 scary music. 565

⁶We computed generalized eta squared statistics (η_G^2) with the aim to yield 566
 measures of effect size comparable across a wide variety of research designs 567
 (Bakeman, 2005), regardless of whether the factor is between or within sub- 568
 jects. These effect-size measures provide indices of effect that are consistent 569
 with Cohen (1988) guidelines indicating that $\eta_G^2 = 0.01$ corresponds to a 570
 small effect, $\eta_G^2 = 0.09$ to a medium effect, and $\eta_G^2 = 0.25$ to a large effect. 571




FIGURE 1 | Mean score and standard error of emotion intensity felt, zygomatic and corrugator muscle activity as a function of intended emotions (happiness, peacefulness, sadness, fear) and age group (younger adults, older adults).

activity, no significant effect of Intended Emotion was found either in older adults, $\chi^2 = 3.00$, $df = 3$, $p = 0.39$, or in younger adults, $\chi^2 = 5.68$, $df = 3$, $p = 0.13$.

INCIDENTAL RECOGNITION TASK

Proportions of hits, false alarms, and corrected recognition scores (hits minus false alarms) are reported in **Table 2**. An analysis of variance was conducted on corrected recognition scores with Age Group (younger adults, older adults) as the between-subjects factor and Intended Emotion (happiness, peacefulness, sadness, fear) as the within-subjects factor. Analysis showed a significant main effect of Age Group indicating that younger adults recognized more musical stimuli than older adults, $F_{(1, 34)} = 30.87$, $p < 0.001$, $\eta_G^2 = 0.50$. There was also a significant main effect of Intended Emotion, $F_{(3, 102)} = 4.86$, $p < 0.05$, $\eta_G^2 = 0.12$, indicating that scary music stimuli were better recognized than peaceful and sad music stimuli. This was confirmed by *post-hoc* Bonferroni comparisons ($ps < 0.05$). No significant Age Group by Intended Emotion interaction was observed, $F_{(3, 102)} = 0.27$, $p = 0.84$, $\eta_G^2 = 0.01$. Younger adults' recognition performances were at chance level or above only for the highly arousing musical stimuli such as happy (52%) and scary (61%) ones, while older adults' recognition performances varied between 10 and 39% through the four intended emotions.

Additional separate analyses conducted on hits and false alarms revealed that older adults generated more false alarms than younger adults, $F_{(1, 34)} = 18.47$, $p < 0.001$, $\eta_G^2 = 0.35$. There was also a significant main effect of Intended Emotion, $F_{(3, 102)} = 22.02$, $p < 0.001$, $\eta_G^2 = 0.36$, as well as a significant Intended Emotion by Age Group interaction for false alarm rates, $F_{(3, 102)} = 4.85$, $p < 0.05$, $\eta_G^2 = 0.08$. *Post-hoc* Bonferroni comparisons showed that the older listeners, compared with their younger counterparts, had more difficulty to correctly reject new happy musical stimuli ($p < 0.05$). Moreover, older adults were better to correctly reject new scary musical stimuli in comparison to all other intended emotions ($ps < 0.001$) while younger adults

were better to correctly reject new scary musical stimuli in comparison to only low arousing stimuli like peaceful ($p < 0.05$) and sad ($p < 0.05$) ones. No significant effect or interaction was found for hits rates. Again, the analysis of covariance with years of education, fluid intelligence scores, and working memory performances as covariates showed that the Intended Emotion by Age Group interaction remained significant, $F_{(3, 93)} = 3.70$, $p < 0.05$, $\eta_G^2 = 0.07$.

D-prime was calculated using tables for d-prime and beta available in Hochhaus (1972) and analyzed using another mixed ANOVA. This indicates the ability to discriminate between true targets and false targets (Green and Swets, 1966), with Age Group (younger adults, older adults) as the between-subjects factor and Intended Emotion (happiness, peacefulness, sadness, fear) as the within-subjects factors. We obtained a significant main effect of Age, $F_{(1, 34)} = 26.41$, $p < 0.001$, $\eta_G^2 = 0.44$ indicating that, overall, older adults showed lower d-prime score ($M = 0.43$, $SE = 0.15$) than younger adults ($M = 1.50$, $SE = 0.15$). This suggests a lower sensitivity in the discrimination of true musical excerpts from false musical excerpts in older adults. No other significant interaction was found with d-prime as the dependent variable. The beta value that indicates the minimum level of activation necessary for a participant to respond to a true target (Green and Swets, 1966) was also calculated (Hochhaus, 1972). No significant main effect or interaction was found with beta value as the dependent variable.

DIFFERENTIATION IN EMOTION FELT

As in previous studies (Vieillard et al., 2008), we derived the best label attributed to each musical excerpt by each participant. This was done selecting the label (i.e., happy, peaceful, sad, scary) that had received the maximal rating. When the maximal rating corresponded to the label that matched the intended emotion, a score of 1 was given. When the maximal rating did not correspond to the emotion, a score of 0 was given. When the highest rating was given for more than one label, the response was considered

Table 2 | Hit rates, False Alarms Rates (Fas) and Corrected Recognition (CR) by Emotions Category and Group Age.

Age group	Intended emotions											
	Happiness			Peacefulness			Sadness			Fear		
	Hits	FA	CR	Hits	FA	CR	Hits	FA	CR	Hits	FA	CR
YOUNG (n = 18)												
M	0.64	0.12	0.52	0.70	0.31	0.39	0.70	0.32	0.38	0.64	0.03	0.61
SE	0.05	0.06	0.05	0.06	0.07	0.07	0.07	0.07	0.09	0.07	0.03	0.07
OLDER (n = 18)												
M	0.74	0.56	0.18	0.66	0.56	0.10	0.63	0.50	0.13	0.46	0.07	0.39
SE	0.05	0.06	0.05	0.06	0.07	0.07	0.07	0.07	0.09	0.07	0.03	0.07

Table 3 | Mean percentage of the label that received the maximal rating of Emotion Felt by younger and older listeners as a function of the Intended Emotions.

Intended Emotion	Emotion felt				
	Happiness	Peacefulness	Sadness	Fear	Ambivalent
Young adults					
Happiness	81	5	2	1	11
Peacefulness	13	37	28	1	21
Sadness	1	20	56	10	13
Fear	10	2	9	72	7
Older adults					
Happiness	73	1	1	1	24
Peacefulness	5	37	22	7	29
Sadness	3	24	35	6	32
Fear	9	4	28	34	25

Bold indicates the match between Emotion Felt and Intended Emotions. Ambivalent responses correspond to highest ratings given to more than one label.

ambivalent and received a score of 0. For example, when an excerpt was perceived eliciting both peacefulness and sadness to the same degree (e.g., with a rating of 7), it was considered as ambivalent. Best labels scores are presented in **Table 3**.

A mixed model analysis of variance was conducted on the mean Best Label with Age Group (younger adults, older adults) as a between-subjects factor and Intended Emotions (happiness, peacefulness, sadness, fear) and Emotion Felt (happiness, peacefulness, sadness, fear) as within-subjects factors. Significant main effects of Age Group, $F_{(1, 34)} = 7.43, p < 0.05, \eta^2_G = 0.18$, and Emotion Felt, $F_{(3, 102)} = 5.87, p < 0.001, \eta^2_G = 0.14$ were found. As expected, results also indicated a significant Intended Emotion by Emotion Felt interaction, $F_{(9, 306)} = 71.45, p < 0.001, \eta^2_G = 0.58$, as well as a significant Intended Emotion by Emotion Felt by Age Group, $F_{(9, 306)} = 5.10, p < 0.001; \eta^2_G = 0.09$. No other significant main effect or interaction was observed. We first compared the experience emotion between younger and older adults for each intended emotion. The results indicated that older adults, compared to younger adults, reported experiencing lower levels of sadness when listening to sad music, $F_{(1, 34)} = 5.44, p < 0.05, \eta^2_G = 0.14$, but reported higher levels of sadness when listening to scary music, $F_{(1, 34)} = 6.35, p < 0.05, \eta^2_G = 0.16$. Moreover,

when listening to scary music, older adults reported experiencing lower levels of fear, $F_{(1, 34)} = 18.78, p < 0.001, \eta^2_G = 0.36$, than their younger counterparts. The second set of comparisons was conducted to compare the emotion experienced for each intended emotion within each age group. The results showed that older adults reported similar levels of sadness and peacefulness when listening to peaceful music, $F_{(1, 34)} = 1.92, p = 0.18, \eta^2_G = 0.05$, and sad music, $F_{(1, 34)} = 0.85, p = 0.36, \eta^2_G = 0.03$, as well as similar levels of sadness and fear when listening to scary music, $F_{(1, 34)} = 0.44, p = 0.51, \eta^2_G = 0.01$. Younger adults reported similar levels of sadness and peacefulness only when listening to peaceful music, $F_{(1, 34)} = 0.73, p = 0.40, \eta^2_G = 0.02$. Happy music was the only music that primarily elicited happiness (when compared with the level of peacefulness felt) in both young and older adults, $F_{(1, 34)} = 117.63, p < 0.001, \eta^2_G = 0.78$, and $F_{(1, 34)} = 105.92, p < 0.001, \eta^2_G = 0.76$, respectively]. The Intended Emotion by Emotion Felt by Age Group interaction remained significant when years of education, fluid intelligence scores, and working memory performances were entered as covariates, $F_{(9, 279)} = 2.90, p < 0.05, \eta^2_G = 0.06$.

THE RELATIONSHIP BETWEEN AGE AND DEPENDENT MEASURES

In order to check for any relationships between age and the different dependent measures (i.e., emotion intensity felt, physiological responses to music, recognition accuracy, and the type of emotion felt) for each of the four intended emotions, we computed a series of correlations. Because age was significantly correlated with years of education, $r_{(34)} = -0.41, p < 0.05$, fluid intelligence, $r_{(34)} = -0.48, p < 0.05$, working memory, $r_{(34)} = -0.35, p < 0.05$, and each of the five measures of auditory thresholds, $r_{(34)} = 0.56, p < 0.001$ for 500 Hz; $r_{(34)} = 0.54, p < 0.001$ for 1000 Hz; $r_{(34)} = 0.61, p < 0.001$ for 2000 Hz; $r_{(34)} = 0.42, p < 0.05$ for 4000 Hz; $r_{(34)} = 0.75, p < 0.001$ for 8000 Hz, these variables were controlled for in partial correlations. Results indicated that the mean scores of hits, corrected recognition, and d-prime for peaceful music were negatively and significantly correlated with age, $r_{(34)} = -0.34, p < 0.05$ for hits, $r_{(34)} = -0.51, p < 0.05$ for corrected recognition, and $r_{(34)} = -0.53, p < 0.05$ for d-prime. Similarly, age was negatively and significantly correlated with the mean score of corrected recognition and d-prime for happy music, $r_{(34)} = -0.69, p < 0.001$ for corrected recognition, and $r_{(34)} = -0.65, p < 0.001$ for d-prime, while it was positively and significantly correlated with the mean score of false alarms, $r_{(34)} = 0.63, p < 0.001$. Altogether, the data indicated that the older the people are, the lower their ability to

799 discriminate studied positive musical excerpts conveying peace-
800 fulness and happiness from unstudied ones. The mean score of
801 Beta index for sad music was also positively and significantly cor-
802 related with age, $r_{(34)} = 0.50$, $p < 0.05$, suggesting that the older
803 the people are, the more conservative they are to discriminate
804 studied stimuli from unstudied musical excerpts conveying sad-
805 ness. Moreover, data indicated that the older the people are, the
806 stronger their experience of sadness while listening to peaceful
807 music, $r_{(34)} = 0.46$, $p < 0.05$, and the weaker their experience of
808 fear while listening to scary music, $r_{(34)} = -0.49$, $p < 0.05$. No
809 other significant correlations were found.

810 **THE RELATIONSHIP BETWEEN EMOTION INTENSITY FELT, FACIAL** 811 **MUSCLE ACTIVITY, AND RECOGNITION PERFORMANCES**

812 For each age group, we investigated to what extent the emotion
813 intensity felt during the first presentation of musical stimuli was
814 linked to the physiological responses as well as to the subsequent
815 cognitive performances on the incidental recognition task. The
816 relationship between physiological reactions and recognition per-
817 formances was also examined. In younger adults, results indicated
818 that the stronger the emotion intensity felt in response to sad
819 music, the higher the hits, $r_{(14)} = 0.54$, $p < 0.05$. In older adults,
820 results showed that the stronger the emotion intensity felt for
821 happy music, the higher the false alarms, $r_{(14)} = 0.54$, $p < 0.05$.
822 No other significant correlations were found.

823 **DISCUSSION**

824 In this study, we investigated how the emotional experience as
825 well as memory recognition for musical excerpts eliciting four
826 different emotions (happiness, sadness, peacefulness, and fear)
827 may change with age. To this end, younger and older listeners
828 were asked to evaluate the intensity of the emotion felt while their
829 facial expressions (i.e., zygomatic and corrugator muscle activity)
830 were recorded. They were then instructed to perform an incidental
831 recognition task followed by another task in which they had
832 to assess for each musical excerpt to what extent they experienced
833 each of the four emotions.

834 As predicted, the results showed that, when presented with
835 happy music, older adults assessed the emotion felt as more
836 intense than their younger counterparts. The fact that older adults
837 rated their emotional experience as significantly more intense
838 for happy music stimuli in comparison to sad and scary music
839 stimuli is consistent with the literature showing that aging is asso-
840 ciated with a relative preference for positivity over negativity. This
841 also supports the view that emotions and motivations cannot
842 be disentangled from each other. However, the assumption that
843 the stronger emotional experience reported by older adults while
844 listening to happy music would be reflected in a greater zygo-
845 matic activity was not supported. Compared to younger adults,
846 older adults showed stronger facial expressions for both corruga-
847 tor and zygomatic muscles as well as for all intended emotions.
848 This suggests that facial expressions are not exclusively aligned
849 with the emotional state, thus raising the question of whether
850 the general increase of facial expressiveness in older adults would
851 simply reflect a deeper engagement in the task. However and
852 interestingly, the current results also showed that older adults'
853 zygomatic activity, but not for young adults', varied as a func-
854 tion of the intended emotions in such a way that older adults

855 showed an increased zygomatic activity for scary excerpts but not
856 for happy or peaceful ones. This is in line with the idea that the
857 reaction of smiling may serve as a defensive goal in inhibiting
858 negative feelings for older adults. It can also be argued that zygo-
859 matic activity may reflect partial facial expression of fear, but then
860 there we should have observed a greater concomitant activity for
861 the corrugator muscle. However, this was not observed. Taken
862 together, these findings are consistent with the idea that older
863 adults' facial expressions possibly reflect an attempt to regulate
864 emotion. Further research is needed to substantiate the role
865 of voluntary facial expressions in the older adults' response to
866 emotions.

867 Consistent with our expectations, our findings indicated that
868 older adults correctly recognized less musical excerpts than their
869 younger counterparts. Moreover, older adults' range of perfor-
870 mances was quite similar to that found by Kensinger (2008) with
871 emotional words. This suggests that modality has little impact on
872 the strength of the memory decline with aging. The results of the
873 present study also indicated that younger adults as well as older
874 adults better recognized negative and arousing musical excerpts
875 (i.e., scary music) than all other excerpts while producing low
876 false alarms rates for these scary music stimuli. This corroborates
877 the hypothesis of an increased distinctiveness of negative stimuli
878 (e.g., Pesta et al., 2001) and extends previous studies that showed
879 that older adults can visually detect arousing and negative stimuli
880 as well as their younger counterparts (e.g., Magai et al., 2006;
881 Knight et al., 2007). Our findings also gave evidence for increased
882 false recognition for happy music stimuli in older adults but
883 not for young adults. We found a negative relationship between
884 age and the ability to discriminate between true and false happy
885 musical excerpts as well as a positive relationship between the
886 emotion intensity felt in older adults and their rate of false
887 alarms for happy stimuli. Taken together, these findings suggest
888 that positive emotion elicited by happy excerpts may produce
889 an attentional bias in older adults that can lead to confusion
890 between studied and non-studied excerpts and thus enhance
891 the probability of false alarms. Such increase in discrimination
892 threshold is consistent with previous studies showing that aging
893 was associated to a higher false response rate to positive words
894 (Fernandes et al., 2008; Piquet et al., 2008) and corroborates the
895 idea that the reduction of distinctiveness for positive information
896 in older adults would be the result of their liberal bias toward
897 positivity. However, in the present study, the positivity bias is
898 detrimental to memory accuracy.

899 Another main finding of the current research is that, when
900 presented with sad and scary musical excerpts, older adults
901 reported experiencing lower levels of sadness and fear than
902 their younger counterparts. Correlation analyses indicated that
903 the older the people are, the weaker their experience of fear
904 felt while listening to scary music. This fits nicely with pre-
905 vious research demonstrating age-related changes in emotion
906 recognition (Laukka and Juslin, 2007; Lima and Castro, 2011)
907 and emotion perception (Vieillard et al., 2012) in music, and
908 extends these studies by showing these changes also occur when
909 participants are focused on their own emotional experience.
910 Interestingly, compared to Lima and Lima and Castro's (2011)
911 recognition paradigm, the personal engagement involved in the
912 current task seems to facilitate the older adults' ability to process

negative emotions. This is consistent with previous findings demonstrating that older adults benefit more from instructions encouraging to focus on emotion than on information acquisition (Mikels et al., 2010) and corroborates the view of an age-related emphasis on emotion processing. Of course, further research is needed to compare the older adults' responsiveness to musical emotions in both contexts of recognition and of emotional experience.

Given the relatively short duration of the musical stimuli used in the present study, one may argue that this could challenge their ability to induce emotions, leading participants to rate their perceived emotions rather than their felt emotions. Although studies aiming to induce felt emotions in listeners tend to use longer excerpts than those investigating perceived emotions (Eerola and Vuoskoski, 2013), we believe that the short excerpts used in our experiment also successfully induced emotions. First of all, our results indicated that participants reported moderate intensity of the emotion felt along with significant differences in facial expressivity. Furthermore, previous findings demonstrated that musical excerpts as short as 13s may recruit neural mechanisms involved in pleasant/unpleasant emotional responses (Blood et al., 1999). In the study of Viellard et al. (2008) which used similar 10 s musical excerpts recorded in a piano timbre, listeners better recognized some intended emotions when focusing on their emotional experience rather than when focusing on the recognition of the emotion. This suggests that asking participants to focus on felt emotions increases the degree of personal engagement in musical emotion even for short musical excerpts. Taken together, these data corroborate the hypothesis that the musical emotions were not only recognized, but indeed felt.

REFERENCES

Allen, R., and Brosgole, L. (1993). Facial and auditory affect recognition in senile geriatrics, the normal elderly and young adults. *Int. J. Neurosci.* 68, 33–42. doi: 10.3109/00207459308994257

Bailey, P. E., Henry, J. D., and Nangle, M. R. (2009). Electromyographic evidence for age-related differences in the mimicry of anger. *Psychol. Aging* 24, 224–229. doi: 10.1037/a0014112

Bakeman, R. (2005). Recommended effect size statistics for repeated measures designs. *Behav. Res. Methods* 37, 379–384. doi: 10.3758/BF03192707

Bayard, S., Erkes, J., and Moroni, C. (2011). Victoria Stroop Test: normative data in a sample group of older people and the study of their clinical applications in the assessment of inhibition in Alzheimer's disease. *Arch. Clin. Neuropsychol.* 26, 653–661. doi: 10.1093/arclin/acr053

Beck, A. T., Steer, R. A., and Brown, G. K. (1998). *Beck Depression Inventory II*. Paris: Center of Applied Psychology.

Blood, A. J., Zatorre, R. J., Bermudez, P., and Evans, A. C. (1999). Emotional responses to pleasant and unpleasant music correlate with activity in paralimbic brain regions. *Nat. Neurosci.* 2, 382. doi: 10.1038/7299

Burriss, L., Powell, D. A., and White, J. (2007). Psychophysiological and subjective indices of emotion as a function of age and gender. *Cogn. Emot.* 21, 182–210. doi: 10.1080/02699930600562235

Cardebat, D., Doyon, B., Puel, M., Goulet, P., and Joannette, Y. (1990). Evocation lexicale formelle et sémantique chez des sujets normaux. Performances et dynamiques de production en fonction du sexe, de l'âge et du niveau d'étude. *Acta Neurol. Belg.* 90, 207–217.

Carstensen, L. L. (2006). The influence of a sense of time on human development. *Science* 312, 1913–1915. doi: 10.1126/science.1127488

Carstensen, L. L., Isaacowitz, D. M., and Charles, S. T. (1999). Taking time seriously. A theory of socioemotional selectivity. *Am. Psychol.* 54, 165–181. doi: 10.1037/0003-066X.54.3.165

One limitation of this study is that we used a cross-sectional design. Historical differences in the cultural system and in musical exposure may have affected young and older adults' performances differently. The observed age differences in emotional responses to music might thus reflect a cohort effect rather than an age effect. Future research would benefit from investigating this issue more thoroughly. Nevertheless, our study suggests that emotional response to music and memory recognition for musical excerpts conveying emotions show differences with advancing age. These age-related differences are characterized by a stronger emotional reactivity for happiness, an increased zygomatic activity in response to scary stimuli, an increase in false recognition for happy musical excerpts, and a decrease in responsiveness to sad and scary music. This study extends previous findings and expands them to music, a powerful channel of emotion communication. Importantly, the findings suggest that aging may cause a decrease in negative affects and an increase in positive affects even when these affects are elicited by a more abstract source of emotion that does not refer to specific events. Finally, the current data are in line with the hypothesis that older adults could use emotional coping skills acquired over the life span in order to avoid potentially negative events and maintain positive ones (Charles et al., 2001; Labouvie-Vief et al., 2010).

ACKNOWLEDGMENTS

This research was funded by the ANR "EMCO" program (Project Strem N ANR 11 EMCO 003 01). We are grateful to Alexandra Richen for her help in running the experiments as well as to Alejandra R. Velasquez and Joanna Blatter-Minn for their proof-reading the manuscript.

Charles, S. T., Mather, M., and Carstensen, L. L. (2003). Aging and emotional memory: the forgettable nature of negative images for older adults. *J. Exp. Psychol. Gen.* 132, 310–324. doi: 10.1037/0096-3445.132.2.310

Charles, S. T., and Piazza, J. R. (2007). Memories of social interactions: age differences in emotional intensity. *Psychol. Aging* 22, 300–309. doi: 10.1037/0882-7974.22.2.300

Charles, S. T., Reynolds, C. A., and Gatz, M. (2001). Age-related differences and change in positive and negative affect over 23 years. *J. Pers. Soc. Psychol.* 80, 136–151. doi: 10.1037/0022-3514.80.1.136

Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences, 2nd Edn*. Hillsdale, NJ: Lawrence Erlbaum Associates Inc.

Drapeau, J., Gosselin, N., Gagnon, L., Peretz, I., and Lorrain, D. (2009). Emotional recognition from face, voice, and music in dementia of the Alzheimer type. *Ann. N.Y. Acad. Sci.* 1169, 342–345. doi: 10.1111/j.1749-6632.2009.04768.x

Eerola, T., and Vuoskoski, J. K. (2013). A review of music and emotion studies: approaches, emotion models, and stimuli. *Music Percept.* 30, 307–340. doi: 10.1525/mp.2012.30.3.307

Fernandes, M., Ross, M., Wiegand, M., and Schryer, E. (2008). Are the memories of older adults positively biased. *Psychol. Aging* 23, 297–306. doi: 10.1037/0882-7974.23.2.297

Gosselin, N., Peretz, I., Johnsen, E., and Adolphs, R. (2007). Amygdala damage impairs emotion recognition from music. *Neuropsychologia* 45, 236–244. doi: 10.1016/j.neuropsychologia.2006.07.012

Gosselin, N., Peretz, I., Noulhiane, M., Hasboun, D., Beckett, C., Baulac, M., et al. (2005). Impaired recognition of scary music following unilateral temporal lobe excision. *Brain* 128, 628–640. doi: 10.1093/brain/awh420

Green, D. M., and Swets, J. A. (1966). Signal detection theory and psychophysics. New York: Wiley.

Grühn, D., Smith, J., and Baltes, P. B. (2005). No aging bias favoring memory for positive material: evidence from a heterogeneity-homogeneity list paradigm

1027 using emotionally toned words. *Psychol. Aging* 20, 579–588. doi: 10.1037/0882-7974.20.4.579

1028 Hochhaus, L. (1972). A table for the

1029 calculation of d' and β . *Psychol. Bull.*

1030 77, 375–376. doi: 10.1037/h0032459

1031 Isaacowitz, D. M., Wadlinger, H. A.,

1032 Goren, D., and Wilson, H. R. (2006a).

1033 Is there an age-related

1034 positivity effect in visual atten-

1035 tion? A comparison of two method-

1036 ologies. *Emotion* 6, 511–516. doi: 10.1037/1528-3542.6.3.511

1037 Isaacowitz, D. M., Wadlinger, H. A.,

1038 Goren, D., and Wilson, H. R. (2006b).

1039 Selective preference in

1040 visual fixation away from negative

1041 images in old age? An eye-tracking

1042 study. *Psychol. Aging* 21, 40–48. doi: 10.1037/0882-7974.21.1.40

1043 Kensinger, E. A. (2008). Age differ-

1044 ences in memory for arousing

1045 and nonarousing emotional

1046 words. *J. Gerontol. B Psychol. Sci. Soc. Sci.* 63, P13–P18. doi: 10.1093/geronb/63.1.P13

1047 Kensinger, E. A., Brierley, B., Medford,

1048 N., Grownold, J. H., and Corkin, S. (2002).

1049 Effects of normal aging and

1050 Alzheimer's disease on emotional

1051 memory. *Emotion* 2, 118–134. doi: 10.1037/1528-3542.2.2.118

1052 Khalfa, S., Peretz, I., Blondin, J.-

1053 P., and Robert, M. (2002). Event-

1054 related skin conductance responses

1055 to musical emotions in humans. *Neurosci. Lett.* 328, 145–149. doi: 10.1016/S0304-3940(02)00462-7

1056 Khalfa, S., Roy, M., Rainville, P., Dalla

1057 Bella, S., and Peretz, I. (2008). Role

1058 of tempo entrainment in

1059 psychophysiological differentiation

1060 of happy and sad music? *Int. J. Psychophysiol.* 68, 17–26. doi: 10.1016/j.ijpsycho.2007.12.001

1061 Knight, M., Seymour, T. L., Gaunt,

1062 J. T., Baker, C., Nesmith, K., and

1063 Mather, M. (2007). Aging and

1064 goal-directed emotional attention:

1065 distraction reverses emotional

1066 biases. *Emotion* 7, 705–714. doi: 10.1037/1528-3542.7.4.705

1067 Labouvie-Vief, G. (2009). Cognition

1068 and equilibrium regulation in devel-

1069 opment and aging. *Restor. Neurol. Neurosci.* 27, 551–565.

1070 Labouvie-Vief, G., Grün, D., and

1071 Studer, J. (2010). "Dynamic inte-

1072 gration of emotion and cognition:

1073 equilibrium regulation in develop-

1074 ment and aging," in *The Handbook of Life-Span Development*, Vol. 2, eds R. M. Lerner, M. E. Lamb, and A. M. Freund (Hoboken, NJ: Wiley), 79–115. doi: 10.1002/9780470880166.hlsd002004

1075 Laukka, P., and Juslin, P. N. (2007).

1076 Similar patterns of age-related dif-

1077 ferences in emotion recognition

1078 from speech and music. *Motiv. Emot.* 31, 182–191. doi: 10.1007/s11031-007-9063-z

1079 Levenson, R. W., Carstensen, L. L.,

1080 Friesen, W. V., and Ekman, P. (1991).

1081 Emotion, physiology, and

1082 expression in old age. *Psychol. Aging* 6, 28–35. doi: 10.1037/0882-7974.6.1.28

1083 Lima, C. F., and Castro, S. L. (2011).

1084 Emotion recognition in music

1085 changes across the adult life span. *Cogn. Emot.* 25, 585–598. doi: 10.1080/02699931.2010.502449

1086 Magai, C., Consedine, N. S.,

1087 Krivoshekova, Y. S., Kudadjie-

1088 Gyamfi, E., and McPherson, R. (2006).

1089 Emotion experience and

1090 expression across the adult life

1091 span: insights from a multimodal

1092 assessment study. *Psychol. Aging* 21, 303–317. doi: 10.1037/0882-7974.21.2.303

1093 Mather, M., Canli, T., English, T.,

1094 Whitfield, S., Wais, P., Ochsner, K.,

1095 et al. (2004). Amygdala responses

1096 to emotionally valenced stimuli in

1097 older and younger adults. *Psychol. Sci.* 15, 259–263. doi: 10.1111/j.0956-7976.2004.00662.x

1098 Mather, M., and Carstensen, L. L. (2005).

1099 Aging and motivated cogni-

1100 tion: the positivity effect in atten-

1101 tion and memory. *Trends Cogn. Sci.* 9, 496–502. doi: 10.1016/j.tics.2005.08.005

1102 Mather, M., and Knight, M. (2005).

1103 Goal-directed memory: the role of

1104 cognitive control in older adults' emotional memory. *Psychol. Aging* 20, 554–570. doi: 10.1037/0882-7974.20.4.554

1105 Mikels, J. A., Larkin, G. R., Reuter-

1106 Lorenz, P. A., and Cartensen, L. L. (2005).

1107 Divergent trajectories in the aging mind: changes in working memory for affective versus visual information with age. *Psychol. Aging* 20, 542–553. doi: 10.1037/0882-7974.20.4.542

1108 Mikels, J. A., Lockenhoff, C. E.,

1109 Maglio, S. J., Goldstein, M. K.,

1110 Garber, A., and Carstensen, L. L. (2010).

1111 Following your heart or your head: focusing on emotions versus information differentially influences the decisions of younger and older adults. *J. Exp. Psychol. Appl.* 16, 87–95. doi: 10.1037/a0018500

1112 Niedenthal, P. M., Barsalou, L. W.,

1113 Winkielman, P., Krauth-Gruber, S., and Ric, F. (2005). Embodiment in attitudes, social perception, and emotion. *Pers. Soc. Psychol. Rev.* 9, 184–211. doi: 10.1207/s15327957pspr0903_1

1114 Pesta, B. J., Murphy, M. D., and Sanders, R. E. (2001). Are emotionally charged lures immune to false memory? *J. Exp. Psychol. Learn. Mem. Cogn.* 27, 328–338. doi: 10.1037/0278-7393.27.2.328

1115 Petit, H., Bakchine, S., Dubois, B.,

1116 Laurent, B., Montagne, B., Touchon, J., et al. (1998). Consensus state-

1117 ment of an interdisciplinary group of French experts on modalities of diagnosis and medical treatment of Alzheimer's disease at a treatable stage. *Rev. Neurol.* 154, 432–438.

1118 Piguet, O., Connally, E., Krendl, A. C., Huot, J. R., and Corkin, S. (2008). False memory in aging: effects of emotional valence on word recognition accuracy. *Psychol. Aging* 23, 307–314. doi: 10.1037/0882-7974.23.2.307

1119 Raven, J., Raven, J. C., and Court, J. H. (1998, updated 2003). *Manual for Raven's Progressive Matrices and Vocabulary Scales. Section 1: General Overview*. San Antonio, TX: Harcourt Assessment.

1120 Reed, A. E., and Carstensen, L. L. (2012). The theory behind the age-related positivity effect. *Front. Psychol.* 3:339. doi: 10.3389/fpsyg.2012.00339

1121 Reitan, R. M. (1958). Validity of the Trail Making Test as an indication of organic brain damage. *Percept. Motor Skills* 8, 271–276. doi: 10.2466/PMS.8.7.271-276

1122 Reminger, S. L., Kaszniak, A. W., and Dalby, P. R. (2000). Age-invariance in the asymmetry of stimulus-evoked emotional facial muscle activity. *Aging Neuropsychol. Cogn.* 7, 156–168. doi: 10.1076/1382-5585(200009)7:3;1-Q:FT156

1123 Scherer, K. R. (2005). What are emotions? And how can they be measured? *Soc. Sci. Inf.* 44, 695–729. doi: 10.1177/0539018405058216

1124 Schneider, W., Eschman, A., and Zuccolotto, A. (2002). *E-Prime User's Guide*. Pittsburgh, PA: Psychology Software Tools Inc.

1125 Smith, D. P., Hillman, C. H., and Duley, A. R. (2005). Influence of age on emotional reactivity during picture processing. *J. Gerontol. B Psychol. Sci. Soc. Sci.* 60, 49–56. doi: 10.1093/geronb/60.1.P49

1126 Spielberger, C. D. (1993). *State and Trait Anxiety Inventory, Form Y*. Paris: Center of Applied Psychology.

1127 Tsai, J. L., Levenson, R. W., and Carstensen, L. L. (2000). Autonomic, subjective, and expressive responses to emotional films in older and younger Chinese Americans and European Americans. *Psychol. Aging* 15, 684–693. doi: 10.1037/0882-7974.15.4.684

1128 van Tricht, M. J., Smeding, H. M., Speelman, J. D., and Schmand, B. A. (2010). Impaired emotion recognition in music in Parkinson's disease. *Brain Cogn.* 74, 58–65. doi: 10.1016/j.bandc.2010.06.005

1129 Viellard, S., Didierjean, A., and Maquestiau, F. (2012). Changes in the perception and the psychological structure of musical emotions with advancing age. *Exp. Aging Res.* 38, 422–441. doi: 10.1080/0361073X.2012.699371

1130 Viellard, S., Peretz, I., Gosselin, N., Khalfa, S., Gagnon, L., and Bouchard, B. (2008). Happy, sad, scary and peaceful musical excerpts for research on emotions. *Cogn. Emot.* 22, 720–752. doi: 10.1080/02699930701503567

1131 Wechsler, D. (2000). *WAIS-III Administration and Scoring Manual*. Paris: Center of Applied Psychology.

1132 Witvliet, C. V. O., and Vrana, S. R. (2007). Play it again Sam: repeated exposure to emotionally evocative music polarises liking and smiling responses, and influences other affective reports, facial EMG, and heart rate. *Cogn. Emot.* 21, 3–25. doi: 10.1080/02699930601000672

1133 Zimmerman, M. E., Brickman, A. M., Paul, R. H., Grieve, S. M., Tate, D. F., Gunstad, J., et al. (2006). The relationship between frontal gray matter volume and cognition varies across the healthy adult lifespan. *Am. J. Geriatr. Psychiatry* 14, 823–833. doi: 10.1097/01.JGP.0000238502.40963.ac

1134

1135 **Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170

1171

1172

1173

1174

1175

1176

1177

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

1250

1251

1252

1253

1254

1255

1256

1257

1258

1259

1260

1261

1262

1263

1264

1265

1266

1267

1268

1269

1270

1271

1272

1273

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

1295

1296

1297

1298

1299

1300

1301

1302

1303

1304

1305

1306

1307

1308

1309

1310

1311

1312

1313

1314

1315

1316

1317

1318

1319

1320

1321

1322

1323

1324

1325

1326

1327

1328

1329

1330

1331

1332

1333

1334

1335

1336

1337

1338

1339

1340

1341

1342

1343

1344

1345

1346

1347

1348

1349

1350

1351

1352

1353

1354

1355

1356

1357

1358

1359

1360

1361

1362

1363

1364

1365

1366

1367

1368

1369

1370

1371

1372

1373

1374

1375

1376

1377

1378

1379

1380

1381

1382

1383

1384

1385

1386

1387

1388

1389

1390

1391

1392

1393

1394

1395

1396

1397

1398

1399

1400

1401

1402

1403

1404

1405

1406

1407

1408

1409

1410

1411

1412

1413

1414

1415

1416

1417

1418

1419

1420

1421

1422

1423

1424

1425

1426

1427

1428

1429

1430

1431

1432

1433

1434

1435

1436

1437

1438

1439

1440

1441

1442

1443

1444

1445

1446

1447

1448

1449

1450

1451

1452

1453

1454

1455

1456

1457

1458

1459

1460

1461

1462

1463

1464

1465

1466

1467

1468

1469

1470

1471

1472

1473

1474

1475

1476

1477

1478

1479

1480

1481

1482

1483

1484

1485

1486

1487

1488

1489

1490

1491

1492

1493

1494

1495

1496

1497

1498

1499

1500